

IN THE CLAIMS

This listing of the claims replaces all prior versions of the claims in the application.

1. (Cancelled)
2. (Cancelled)
3. (Currently Amended) An isolated polynucleotide encoding a polypeptide ~~of claim 1~~
selected from the group consisting of:
 - a) a polypeptide comprising the amino acid sequence of SEQ ID NO:1,
 - b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to the amino acid sequence of SEQ ID NO:1, and
 - c) an immunogenic fragment of at least 10 contiguous amino acids of a polypeptide having the amino acid sequence of SEQ ID NO:1.
4. (Currently Amended) An isolated polynucleotide ~~encoding a polypeptide~~ of claim 2
which encodes a polypeptide comprising the amino acid sequence of SEQ ID NO:1.
5. (Original) An isolated polynucleotide of claim 4 comprising the polynucleotide sequence of SEQ ID NO:2.
6. (Original) A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 3.
7. (Original) A cell transformed with a recombinant polynucleotide of claim 6.
8. (Canceled)

9. (Currently Amended) A method of producing a polypeptide ~~of claim 1~~ encoded by a polynucleotide of claim 3, the method comprising:

- a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide ~~encoding the polypeptide of claim 1~~ of claim 3, and
- b) recovering the polypeptide so expressed.

10. (Original) A method of claim 9, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO:1.

11. (Cancelled)

12. (Original) An isolated polynucleotide selected from the group consisting of:

- a) a polynucleotide comprising the polynucleotide sequence of SEQ ID NO:2,
- b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to the polynucleotide sequence of SEQ ID NO:2,
- c) a polynucleotide complementary to a polynucleotide of a),
- d) a polynucleotide complementary to a polynucleotide of b), and
- e) an RNA equivalent of a)-d).

13. (Original) An isolated polynucleotide comprising at least 60 contiguous nucleotides of a polynucleotide of claim 12.

14. (Original) A method of detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 12, the method comprising:

- a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and
- b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.

15. (Original) A method of claim 14, wherein the probe comprises at least 60 contiguous nucleotides.

16. (Original) A method of detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 12, the method comprising:

- a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and
- b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

Claims 17-27. (Cancelled)

28. (Original) A method of screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 5, the method comprising:

- a) exposing a sample comprising the target polynucleotide to a compound, under conditions suitable for the expression of the target polynucleotide,
- b) detecting altered expression of the target polynucleotide, and
- c) comparing the expression of the target polynucleotide in the presence of varying amounts of the compound and in the absence of the compound.

29. (Original) A method of assessing toxicity of a test compound, the method comprising:

- a) treating a biological sample containing nucleic acids with the test compound,
- b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide of claim 12 under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide comprising a polynucleotide sequence of a polynucleotide of claim 12 or fragment thereof,
- c) quantifying the amount of hybridization complex, and
- d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

Claims 30-55. (Cancelled)